

KRYSTEXXA REFERRAL FORM

PATIENT DEMOGRAPHICS:		
PATIENT NAME:	PATIENT'S CONTACT #:	
DATE OF REFERRAL:	ADDRESS:	
DATE OF BIRTH:	CITY, STATE, ZIP:	
FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:		
HEIGHT: _____ FEET _____ INCHES	GENDER: FEMALE MALE	
WEIGHT: _____ LB or _____ KG	ALLERGIES: SEE LIST NKDA	
PRIMARY DIAGNOSIS:		
M1A.9XX0 - Chronic Gout, without Tophus (Tophi) M1A.9XX1 - Chronic Gout, with Tophus (Tophi)		
Other: _____		
REQUIRED DOCUMENTATION: Please provide a copy of the following documents.		
<input checked="" type="checkbox"/> INSURANCE CARD (Front & Back) <input checked="" type="checkbox"/> PATIENT DEMOGRAPHICS <input checked="" type="checkbox"/> MOST RECENT LABS <input checked="" type="checkbox"/> MEDICATION LIST <input checked="" type="checkbox"/> H&P		
<input checked="" type="checkbox"/> Prior Failed or Intolerant Gout Therapy (if any): Allopurinol Febuxostat Probenecid Other: _____		
<input checked="" type="checkbox"/> Serum Uric Acid Level: _____ Date Drawn: _____		
<input checked="" type="checkbox"/> G6PD Results: _____ Date Drawn: _____ -OR- G6PD to be drawn by FlexCare		
PRIMARY MEDICATION ORDER:		
Krystexxa 8 mg IV every 2 weeks Other: _____		
*Manufacturer recommendations now include co-administration with Methotrexate 15 mg PO weekly and Folic Acid 1 mg PO daily beginning 4 weeks prior to initiation of Krystexxa. Immunomodulation to be prescribed/managed by FlexCare providers		
Labs to be drawn by FlexCare: <input checked="" type="checkbox"/> Uric Acid (within 48 hrs prior to each infusion)		
Lab Frequency: _____		
First Dose: Y N		
<input checked="" type="checkbox"/> Refill x12 months unless otherwise noted		
PRN & PREMEDICATIONS:		
	30 minutes prior to every infusion	PRN
MEDICATIONS		
Acetaminophen 650 mg PO		PRN every _____ hours for mild or moderate infusion reaction.
Diphenhydramine 25 mg PO		PRN every _____ hours for mild or moderate infusion reaction.
Diphenhydramine 25 mg IV	<input checked="" type="checkbox"/>	PRN every _____ hours for mild or moderate infusion reaction.
Methylprednisolone 125 mg IV	<input checked="" type="checkbox"/>	PRN every _____ hours for mild or moderate infusion reaction.
Other: _____		PRN every _____ hours for mild or moderate infusion reaction.
LINE USE/CARE ORDERS:		
<input checked="" type="checkbox"/> START PIV/ACCESS CVC <input checked="" type="checkbox"/> FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (SEE REVERSE SIDE) OTHER FLUSH ORDERS: (please fax other orders if checking this box)		
ADVERSE REACTION & ANAPHYLAXIS ORDERS:		
<input checked="" type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side) OTHER: (please fax other reaction orders if checking this box)		
PRESCRIBER INFORMATION: Please check preferred form of communication.		
PROVIDER NAME:	PHONE:	
OFFICE CONTACT:	FAX:	
ADDRESS:	EMAIL:	
CITY, STATE, ZIP:	NPI:	
PROVIDER SIGNATURE: DATE:		

KRYSTEXXA REFERRAL FORM

FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

**This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INFUSION REACTION	MODERATE INFUSION REACTION	SEVERE INFUSION REACTION/ANAPHYLAXIS
SYMPTOM CLASSIFICATION	<ul style="list-style-type: none"> Flushing Dizziness Headache Apprehension Diaphoresis Palpitations Nausea / Vomiting Pruritis 	<ul style="list-style-type: none"> Chest Tightness Shortness of Breath Hypo/hypertension (>20 mmHg Change in Systolic BP from Baseline) Increased Temperature (>2 Degrees Fahrenheit) Urticaria 	<ul style="list-style-type: none"> Hypo/hypertension (>40 mmHg Change in Systolic BP from Baseline). Increase Temperature (>2 Degrees Fahrenheit) with Rigors Shortness of Breath with Wheezing Laryngeal Edema Chest Pain Hypoxemia
TREATMENT PROTOCOL FOR ADULTS >66LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

FLUSHING PROTOCOLS

		FLUSHING PROTOCOL Normal Saline*		LOCKING PROTOCOL Heparin Sodium	
		0.9% Sodium Chloride		10 Units/mL	100 Units/mL
PATIENT CLASSIFICATION	LINE TYPE	PRE-ADMIN	POST ADMIN	POST LAB DRAW	POST NS FLUSH*
ADULT > 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	10 mL	10 mL	5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL
	Tunneled & non-Tunneled Catheters	5 mL	10 mL	10 mL	5 mL
PEDIATRIC 33 LBS - 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	5 mL	10 mL	3 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	5 mL	10 mL	3 mL
	Tunneled & non-Tunneled Catheters	5 mL	5 mL	10 mL	3 mL

FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.