

## REMICADE REFERRAL FORM

### PATIENT DEMOGRAPHICS:

PATIENT NAME:	PATIENT'S CONTACT #:
DATE OF REFERRAL:	ADDRESS:
DATE OF BIRTH:	CITY, STATE, ZIP:
FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:	
HEIGHT: _____ FEET _____ INCHES	GENDER: FEMALE MALE
WEIGHT: _____ LB or _____ KG	ALLERGIES: SEE LIST NKDA

### PRIMARY DIAGNOSIS:

L40.5 - Arthropathic psoriasis	M45 - Ankylosing spondylitis	K50 - Crohn's disease (regional enteritis)
K50.01 - Crohn's disease of small intestine w/ comp	K50.11 - Crohn's disease of large intestine w/ comp	K50.81 - Crohn's disease, small & large intestine w/ comp
K51 - Ulcerative Colitis	K50.91 - Crohn's disease, unspecified, w/ comp	K51.01 - Ulcerative (chronic) pancolitis w/ comp
K51.21 - Ulcerative (chronic) proctitis w/ comp	K51.31 - Ulcerative (chronic) rectosigmoiditis w/ comp	K51.51 - Left sided colitis w/ comp
K51.81 - Other ulcerative colitis w/ comp	K51.91 - Ulcerative colitis, unspecified, w/ comp	M05.7XX - RA w/ rheum factor w/o organ/systems involvement
M05.8XX - Other rheumatoid arthritis w/ rheum factor	M05.9 - Rheumatoid arthritis w/ rheum factor, unspec	M06 - Other Rheumatoid arthritis
M06.0XX - Rheumatoid arthritis w/o rheum factor	M06.8XX - Other specified rheumatoid arthritis	M06.9 - Rheumatoid arthritis, unspecified
Other: _____		

### REQUIRED DOCUMENTATION: Please provide a copy of the following documents.

<input checked="" type="checkbox"/> INSURANCE CARD (front & back)	<input checked="" type="checkbox"/> PATIENT DEMOGRAPHICS	<input checked="" type="checkbox"/> MOST RECENT LABS	<input checked="" type="checkbox"/> MEDICATION LIST
<input checked="" type="checkbox"/> HISTORY & PHYSICAL	<input checked="" type="checkbox"/> TRIED/FAILED THERAPIES	<input checked="" type="checkbox"/> NEGATIVE TB RESULTS	

### PRIMARY MEDICATION ORDER:

Remicade 3 mg/kg IV at weeks 0, 2, 6, and every \_\_\_\_\_ weeks thereafter.

Remicade 5 mg/kg IV at weeks 0, 2, 6, and every \_\_\_\_\_ weeks thereafter.

Remicade 8 mg/kg IV at weeks 0, 2, 6, and every \_\_\_\_\_ weeks thereafter.

Remicade 10 mg/kg IV at weeks 0, 2, 6, and every \_\_\_\_\_ weeks thereafter.

Remicade \_\_\_\_\_ mg/kg IV every \_\_\_\_\_ weeks.

Other: \_\_\_\_\_

FIRST DOSE: Y N

Biosimilar may be used according to payer guidelines, unless otherwise noted

Refill x12 months unless otherwise noted

### PRN & PREMEDICATIONS:

MEDICATIONS	30 minutes prior to every infusion	PRN
Acetaminophen 650 mg PO		PRN every _____ hours for mild or moderate infusion reaction.
Diphenhydramine 25 mg PO		PRN every _____ hours for mild or moderate infusion reaction.
Diphenhydramine 25 mg IV		PRN every _____ hours for mild or moderate infusion reaction.
Methylprednisolone 125 mg IV		PRN every _____ hours for mild or moderate infusion reaction.
Other: _____		PRN every _____ hours for mild or moderate infusion reaction.

### LINE USE/CARE ORDERS:

START PIV/ACCESS CVC

FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (SEE REVERSE SIDE)

OTHER FLUSH ORDERS: (please fax other reaction orders if checking this box)

### ADVERSE REACTION & ANAPHYLAXIS ORDERS:

ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side)

OTHER: (please fax other reaction orders if checking this box)

### LAB ORDERS: Please include frequency.


### PRESCRIBER INFORMATION: Please check preferred form of communication.

PROVIDER NAME:	PHONE:
OFFICE CONTACT:	FAX:
ADDRESS:	EMAIL:
CITY, STATE, ZIP:	NPI:

→  
PROVIDER SIGNATURE:

DATE:

# REMICADE REFERRAL FORM

## FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

*\*This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INFUSION REACTION	MODERATE INFUSION REACTION	SEVERE INFUSION REACTION/ANAPHYLAXIS
<b>SYMPTOM CLASSIFICATION</b>	<ul style="list-style-type: none"> <li>Flushing</li> <li>Dizziness</li> <li>Headache</li> <li>Apprehension</li> <li>Diaphoresis</li> <li>Palpitations</li> <li>Nausea / Vomiting</li> <li>Pruritis</li> </ul>	<ul style="list-style-type: none"> <li>Chest Tightness</li> <li>Shortness of Breath</li> <li>Hypo/hypertension (&gt;20 mmHg Change in Systolic BP from Baseline)</li> <li>Increased Temperature (&gt;2 Degrees Fahrenheit)</li> <li>Urticaria</li> </ul>	<ul style="list-style-type: none"> <li>Hypo/hypertension (&gt;40 mmHg Change in Systolic BP from Baseline).</li> <li>Increase Temperature (&gt;2 Degrees Fahrenheit) with Rigors</li> <li>Shortness of Breath with Wheezing</li> <li>Laryngeal Edema</li> <li>Chest Pain</li> <li>Hypoxemia</li> </ul>
<b>TREATMENT PROTOCOL FOR ADULTS &gt;66LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
<b>TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

**FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

## FLUSHING PROTOCOLS

		FLUSHING PROTOCOL Normal Saline*		LOCKING PROTOCOL Heparin Sodium	
		0.9% Sodium Chloride		10 Units/mL	100 Units/mL
PATIENT CLASSIFICATION	LINE TYPE	PRE-ADMIN	POST ADMIN	POST LAB DRAW	POST NS FLUSH*
<b>ADULT &gt; 66 LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	10 mL	10 mL	5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL
	Tunneled & non-Tunneled Catheters	5 mL	10 mL	10 mL	5 mL
<b>PEDIATRIC 33 LBS - 66 LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	5 mL	10 mL	3 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	5 mL	10 mL	3 mL
	Tunneled & non-Tunneled Catheters	5 mL	5 mL	10 mL	3 mL

**FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

\*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.