

## TEZSPIRE REFERRAL FORM

### PATIENT DEMOGRAPHICS:

PATIENT NAME:	PATIENT'S CONTACT #:		
DATE OF REFERRAL:	ADDRESS:		
DATE OF BIRTH:	CITY, STATE, ZIP:		
FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:			
HEIGHT: _____ FEET _____ INCHES	GENDER: FEMALE MALE		
WEIGHT: _____ LB or _____ KG	ALLERGIES: SEE LIST NKDA		

### PRIMARY DIAGNOSIS:

J45.50 Severe persistent asthma, uncomplicated	J45.51 Severe persistent asthma with (acute) exacerbation
Other	

### REQUIRED DOCUMENTATION: Please provide a copy of the following documents.

1. INSURANCE CARD (Front & Back)  
  2. PATIENT DEMOGRAPHICS  
  3. MOST RECENT LABS  
  4. MEDICATION LIST  
 5. H & P  
  6. TRIED/FAILED THERAPIES

### PRIMARY MEDICATION ORDER:

### PRN & PREMEDICATIONS:

Please include MEDICATION, DOSE, FREQUENCY, DURATION and any ADDITIONAL administration INSTRUCTIONS specific to the primary therapy.  Tezspire 210 mg subcutaneous injection every 4 weeks.  Other: _____ _____  FIRST DOSE:      Y            N <input checked="" type="checkbox"/> Refill x12 months unless otherwise noted.	MEDICATIONS	60 minutes prior to every injection	PRN
	Acetaminophen 650 mg PO		PRN every ___ hour for mild or moderate injection reaction.
	Diphenhydramine 50 mg PO		PRN every ___ hour for mild or moderate injection reaction.
	Diphenhydramine 25 mg IV		PRN every ___ hour for mild or moderate injection reaction.
	Methylprednisolone 125 mg IV		PRN every ___ hour for mild or moderate injection reaction.
	Other: _____		PRN every ___ hour for mild or moderate injection reaction.

### ADVERSE REACTION & ANAPHYLAXIS ORDERS:

ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side)  
 START PIV/ACCESS CVC (As Needed for ADVERSE REACTIONS)  
 OTHER: (please fax other reaction orders if checking this box)

### PRESCRIBER INFORMATION: Please check preferred form of communication.

PROVIDER NAME:	PHONE:
OFFICE CONTACT:	FAX:
ADDRESS:	EMAIL:
CITY, STATE, ZIP:	NPI:
_____ → (GENERIC SUBSTITUTION PERMITTED) PROVIDER SIGNATURE: _____ DATE: _____	
_____ → (DISPENSE AS WRITTEN) PROVIDER SIGNATURE: _____ DATE: _____	

# TEZSPIRE REFERRAL FORM

## FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

*\*This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INJECTION REACTION	MODERATE INJECTION REACTION	SEVERE INJECTION REACTION/ANAPHYLAXIS
<b>SYMPTOM CLASSIFICATION</b>	<ul style="list-style-type: none"> <li>Flushing</li> <li>Dizziness</li> <li>Headache</li> <li>Apprehension</li> <li>Diaphoresis</li> <li>Palpitations</li> <li>Nausea / Vomiting</li> <li>Pruitis</li> </ul>	<ul style="list-style-type: none"> <li>Chest Tightness</li> <li>Shortness of Breath</li> <li>Hypo/hypertension (&gt;20 mmHg Change in Systolic BP from Baseline)</li> <li>Increased Temperature (&gt;2 Degrees Fahrenheit)</li> <li>Urticaria</li> </ul>	<ul style="list-style-type: none"> <li>Hypo/hypertension (&gt;40 mmHg Change in Systolic BP from Baseline).</li> <li>Increase Temperature (&gt;2 Degrees Fahrenheit) with Rigors</li> <li>Shortness of Breath with Wheezing</li> <li>Laryngeal Edema</li> <li>Chest Pain</li> <li>Hypoxemia</li> </ul>
<b>TREATMENT PROTOCOL FOR ADULTS &gt;66LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
<b>TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS</b>	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% naCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

**FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

## FLUSHING PROTOCOLS

		FLUSHING PROTOCOL Normal Saline*		LOCKING PROTOCOL Heparin Sodium	
		0.9% Sodium Chloride		10 Units/mL	100 Units/mL
PATIENT CLASSIFICATION	LINE TYPE	PRE-ADMIN	POST ADMIN	POST LAB DRAW	POST NS FLUSH*
<b>ADULT &gt; 66 LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	10 mL	10 mL	5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL
	Tunneled & non-Tunneled Catheters	5 mL	10 mL	10 mL	5 mL
<b>PEDIATRIC 33 LBS - 66 LBS</b>	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	5 mL	10 mL	3 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	5 mL	10 mL	3 mL
	Tunneled & non-Tunneled Catheters	5 mL	5 mL	10 mL	3 mL

**FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.**

\*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.