

LUPUS REFERRAL FORM

PATIENT DEMOGRAPHICS:

PATIENT NAME:	PATIENT'S CONTACT #:	
DATE OF REFERRAL:	ADDRESS:	
DATE OF BIRTH:	CITY, STATE, ZIP:	
FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:		
HEIGHT: _____ FEET _____ INCHES	GENDER: FEMALE	MALE
WEIGHT: _____ LB or _____ KG	ALLERGIES: SEE LIST	NKDA

PRIMARY DIAGNOSIS:

M32.10 - Systemic lupus erythematosus, organ or system involvement unspecified	M32.19 - Other organ or system involvement in systemic lupus erythematosus
M32.8 - Other forms of systemic lupus erythematosus	M32.9 - Systemic lupus erythematosus, unspecified
M32.14 - Glomerular disease in systemic lupus erythematosus	M32.15 - Tubulo-interstitial nephropathy in systemic lupus erythematosus

REQUIRED DOCUMENTATION: Please provide a copy of the following documents.

1. INSURANCE CARD (Front & Back)
 2. PATIENT DEMOGRAPHICS
 3. MOST RECENT LABS
 4. MEDICATION LIST
 5. H & P
 6. TRIED/FAILED THERAPIES

PRIMARY MEDICATION ORDER:

Benlysta
Initial: 10 mg/kg IV at weeks 0, 2, 4, and every 4 weeks thereafter.
Maintenance: 10 mg/kg IV every 4 weeks.

Saphnelo
300 mg IV every 4 weeks.

Other: _____

FIRST DOSE: Y N

Refill x12 months unless otherwise noted.

PRN & PREMEDICATIONS:

MEDICATIONS	30 minutes prior to every infusion	PRN
Acetaminophen 650 mg PO		PRN every ___ hour for mild or moderate infusion reaction.
Diphenhydramine 25 mg PO		PRN every ___ hour for mild or moderate infusion reaction.
Diphenhydramine 25 mg IV		PRN every ___ hour for mild or moderate infusion reaction.
Methylprednisolone 125 mg IV		PRN every ___ hour for mild or moderate infusion reaction.
Other: _____		PRN every ___ hour for mild or moderate infusion reaction.

LINE USE/CARE ORDERS:

START PIV/ACCESS CVC
 FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (SEE REVERSE SIDE)
 OTHER FLUSH ORDERS: (please fax other reaction orders if checking this box)

ADVERSE REACTION & ANAPHYLAXIS ORDERS:

ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side)
 OTHER: (please fax other reaction orders if checking this box)

PRESCRIBER INFORMATION: Please check preferred form of communication.

PHYSICIAN NAME:	PHONE:
OFFICE CONTACT:	FAX:
ADDRESS:	EMAIL:
CITY, STATE, ZIP:	NPI:

 (GENERIC SUBSTITUTION PERMITTED)
 PHYSICIAN SIGNATURE:

DATE:

 (DISPENSE AS WRITTEN)
 PHYSICIAN SIGNATURE:

DATE:

LUPUS REFERRAL FORM

FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

**This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INFUSION REACTION	MODERATE INFUSION REACTION	SEVERE INFUSION REACTION/ANAPHYLAXIS
SYMPTOM CLASSIFICATION	<ul style="list-style-type: none"> Flushing Dizziness Headache Apprehension Diaphoresis Palpitations Nausea / Vomiting Pruitis 	<ul style="list-style-type: none"> Chest Tightness Shortness of Breath Hypo/hypertension (>20 mmHg Change in Systolic BP from Baseline) Increased Temperature (>2 Degrees Fahrenheit) Urticaria 	<ul style="list-style-type: none"> Hypo/hypertension (>40 mmHg Change in Systolic BP from Baseline). Increase Temperature (>2 Degrees Fahrenheit) with Rigors Shortness of Breath with Wheezing Laryngeal Edema Chest Pain Hypoxemia
TREATMENT PROTOCOL FOR ADULTS >66LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

FOR CHILDREN < 33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

FLUSHING PROTOCOLS

		FLUSHING PROTOCOL Normal Saline*		LOCKING PROTOCOL Heparin Sodium	
		0.9% Sodium Chloride		10 Units/mL	100 Units/mL
PATIENT CLASSIFICATION	LINE TYPE	PRE-ADMIN	POST ADMIN	POST LAB DRAW	POST NS FLUSH*
ADULT > 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	10 mL	10 mL	5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL
	Tunneled & non-Tunneled Catheters	5 mL	10 mL	10 mL	5 mL
PEDIATRIC 33 LBS - 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	5 mL	10 mL	3 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	5 mL	10 mL	3 mL
	Tunneled & non-Tunneled Catheters	5 mL	5 mL	10 mL	3 mL

FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.