



UPLIZNA REFERRAL FORM

PATIENT DEMOGRAPHICS:																					
PATIENT NAME:	PATIENT'S CONTACT #:																				
DATE OF REFERRAL:	ADDRESS:																				
DATE OF BIRTH:	CITY, STATE, ZIP:																				
FOR COPAY ASSISTANCE, PLEASE PROVIDE LAST 4 OF SSN:																					
HEIGHT: _____ FEET _____ INCHES	GENDER: FEMALE MALE																				
WEIGHT: _____ LB or _____ KG	ALLERGIES: SEE LIST NKDA																				
PRIMARY DIAGNOSIS:																					
G36.0 - Neuromyelitis optica [Devic]																					
Other: _____																					
REQUIRED DOCUMENTATION: Please provide a copy of the following documents.																					
<input checked="" type="checkbox"/> 1. INSURANCE CARD (Front & Back) <input checked="" type="checkbox"/> 2. PATIENT DEMOGRAPHICS <input checked="" type="checkbox"/> 3. MOST RECENT LABS <input checked="" type="checkbox"/> 4. MEDICATION LIST <input checked="" type="checkbox"/> 5. H & P <input checked="" type="checkbox"/> 6. POSITIVE LAB RESULTS FOR ANTI-AQUAPORIN-4 (AQP4) ANTIBODIES <input checked="" type="checkbox"/> 7. DOCUMENTATION OF CLINICAL SIGNS AND SYMPTOMS (EX. EDSS SCORE, ATTACK HISTORY) <input checked="" type="checkbox"/> 8. PREVIOUS TRIED/FAILED THERAPIES WITH CONTRAINDICATIONS <input checked="" type="checkbox"/> 9. HEP B, TB, QUANTITATIVE SERUM IMMUNOGLOBULINS																					
PRIMARY MEDICATION ORDER:																					
<p>Initial – Uplizna 300 mg IV at day 1 and day 15.</p> <p>Maintenance Only – Uplizna 300 mg IV every 6 months (starting 6 months from the 1st infusion).</p> <p>Other: _____</p> <p><input checked="" type="checkbox"/> Labs Orders (to be drawn at infusion clinic): _____ with each dose every _____ weeks</p> <p>FIRST DOSE: Y N</p> <p><input checked="" type="checkbox"/> Refill x12 months unless otherwise noted.</p>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #cccccc;"> <th colspan="2">PRN & PREMEDICATIONS:</th> </tr> <tr> <th style="width: 50%;">MEDICATIONS</th> <th style="width: 20%;">30 minutes prior to every infusion</th> <th style="width: 30%;">PRN</th> </tr> </thead> <tbody> <tr> <td>Acetaminophen 650 mg PO</td> <td style="text-align: center;"><input checked="" type="checkbox"/></td> <td>PRN every ____ hour for mild or moderate infusion reaction.</td> </tr> <tr> <td>Diphenhydramine 25 mg PO</td> <td style="text-align: center;"><input checked="" type="checkbox"/></td> <td>PRN every ____ hour for mild or moderate infusion reaction.</td> </tr> <tr> <td>Diphenhydramine 25 mg IV</td> <td></td> <td>PRN every ____ hour for mild or moderate infusion reaction.</td> </tr> <tr> <td>Methylprednisolone 125 mg IV</td> <td style="text-align: center;"><input checked="" type="checkbox"/></td> <td>PRN every ____ hour for mild or moderate infusion reaction.</td> </tr> <tr> <td>Other: _____</td> <td></td> <td>PRN every ____ hour for mild or moderate infusion reaction.</td> </tr> </tbody> </table>	PRN & PREMEDICATIONS:		MEDICATIONS	30 minutes prior to every infusion	PRN	Acetaminophen 650 mg PO	<input checked="" type="checkbox"/>	PRN every ____ hour for mild or moderate infusion reaction.	Diphenhydramine 25 mg PO	<input checked="" type="checkbox"/>	PRN every ____ hour for mild or moderate infusion reaction.	Diphenhydramine 25 mg IV		PRN every ____ hour for mild or moderate infusion reaction.	Methylprednisolone 125 mg IV	<input checked="" type="checkbox"/>	PRN every ____ hour for mild or moderate infusion reaction.	Other: _____		PRN every ____ hour for mild or moderate infusion reaction.
PRN & PREMEDICATIONS:																					
MEDICATIONS	30 minutes prior to every infusion	PRN																			
Acetaminophen 650 mg PO	<input checked="" type="checkbox"/>	PRN every ____ hour for mild or moderate infusion reaction.																			
Diphenhydramine 25 mg PO	<input checked="" type="checkbox"/>	PRN every ____ hour for mild or moderate infusion reaction.																			
Diphenhydramine 25 mg IV		PRN every ____ hour for mild or moderate infusion reaction.																			
Methylprednisolone 125 mg IV	<input checked="" type="checkbox"/>	PRN every ____ hour for mild or moderate infusion reaction.																			
Other: _____		PRN every ____ hour for mild or moderate infusion reaction.																			
LINE USE/CARE ORDERS:																					
<p><input checked="" type="checkbox"/> START PIV/ACCESS CVC</p> <p><input checked="" type="checkbox"/> FLUSH DEVICE PER FLEXCARE INFUSION POLICY & PROCEDURE (SEE REVERSE SIDE)</p> <p>OTHER FLUSH ORDERS: (please fax other reaction orders if checking this box)</p>	<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr style="background-color: #cccccc;"> <th>ADVERSE REACTION & ANAPHYLAXIS ORDERS:</th> </tr> </thead> <tbody> <tr> <td> <input checked="" type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side) OTHER: (please fax other reaction orders if checking this box) </td> </tr> </tbody> </table>	ADVERSE REACTION & ANAPHYLAXIS ORDERS:	<input checked="" type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side) OTHER: (please fax other reaction orders if checking this box)																		
ADVERSE REACTION & ANAPHYLAXIS ORDERS:																					
<input checked="" type="checkbox"/> ADMINISTER ACUTE INFUSION AND ANAPHYLAXIS MEDICATIONS PER FLEXCARE INFUSION POLICY AND PROCEDURE (See Reverse Side) OTHER: (please fax other reaction orders if checking this box)																					
PRESCRIBER INFORMATION: Please check preferred form of communication.																					
PROVIDER NAME:	PHONE:																				
OFFICE CONTACT:	FAX:																				
ADDRESS:	EMAIL:																				
CITY, STATE, ZIP:	NPI:																				
PROVIDER SIGNATURE: _____																					
DATE: _____																					

UPLIZNA REFERRAL FORM

FLEXCARE INFUSION CENTER'S ACUTE & ANAPHYLAXIS MEDICATION PROTOCOL:

**This table does not reflect non-medicinal interventions that are part of FlexCare's protocol, such as slowing or stopping the infusion and physician/911 notification.*

	MILD INFUSION REACTION	MODERATE INFUSION REACTION	SEVERE INFUSION REACTION/ANAPHYLAXIS
SYMPTOM CLASSIFICATION	<ul style="list-style-type: none"> Flushing Dizziness Headache Apprehension Diaphoresis Palpitations Nausea / Vomiting Pruitis 	<ul style="list-style-type: none"> Chest Tightness Shortness of Breath Hypo/hypertension (>20 mmHg Change in Systolic BP from Baseline) Increased Temperature (>2 Degrees Fahrenheit) Urticaria 	<ul style="list-style-type: none"> Hypo/hypertension (>40 mmHg Change in Systolic BP from Baseline). Increase Temperature (>2 Degrees Fahrenheit) with Rigors Shortness of Breath with Wheezing Laryngeal Edema Chest Pain Hypoxemia
TREATMENT PROTOCOL FOR ADULTS >66LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500 mL at 125mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 50 mg IV or IM Inject epinephrine 0.3mg/0.3 mL IM into the midanterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl 1000mL bolus for an incomplete response to IM epinephrine. May repeat x1.
TREATMENT PROTOCOL FOR CHILDREN 33LBS - 66 LBS	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Administer PRN medications per Physician order	<input checked="" type="checkbox"/> Apply oxygen via ambu bag or high flow nasal canula, if vomiting. <input checked="" type="checkbox"/> Administer 0.9% NaCl 500mL at 75mL/hr to maintain IV access. <input checked="" type="checkbox"/> Administer diphenhydramine 1-2 mg/kg IM or slow IVP not to exceed 25mg/min <input checked="" type="checkbox"/> Inject epinephrine 0.15mg/0.15 mL IM into the mid-anterolateral aspect of the thigh; repeat in 5-15 minutes if needed. <input checked="" type="checkbox"/> Administer 0.9% NaCl bolus 20mL/kg for an incomplete response to IM epinephrine. May repeat x1.

FOR CHILDREN <33 LBS FLEXCARE INFUSION UTILIZES THE REACTION ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

FLUSHING PROTOCOLS

		FLUSHING PROTOCOL Normal Saline*		LOCKING PROTOCOL Heparin Sodium	
		0.9% Sodium Chloride		10 Units/mL	100 Units/mL
PATIENT CLASSIFICATION	LINE TYPE	PRE-ADMIN	POST ADMIN	POST LAB DRAW	POST NS FLUSH*
ADULT > 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	10 mL	10 mL	5 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	10 mL	10 mL	5 mL
	Tunneled & non-Tunneled Catheters	5 mL	10 mL	10 mL	5 mL
PEDIATRIC 33 LBS - 66 LBS	Peripheral IV Catheter	3 mL	3 mL		3 mL
	Midline	3 mL	3 mL		3 mL
	Implanted Port	5 mL	5 mL	10 mL	3 mL
	Peripherally Inserted Central Catheters (PICC)	5 mL	5 mL	10 mL	3 mL
	Tunneled & non-Tunneled Catheters	5 mL	5 mL	10 mL	3 mL

FOR CHILDREN <33 LBS, FLEXCARE INFUSION UTILIZES THE FLUSHING ORDERS OBTAINED BY THE REFERRING PHYSICIAN.

*0.9% NS will be substituted with Dextrose 5% or alternative only when indicated due to medication incompatibility with NS.